

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry**Exercise 26A**

Page No: 784

Question 1: If a point lies on the z-axis, then find its x-coordinate and y-coordinate.**Solution:**

If the point lies on the z axis then its x-coordinate and y-coordinate will be zero.

Question 2: If a point lies on yz-plane then what is its x-coordinate?**Solution:**

If the point lies on yz-plane then x-coordinate will be zero.

Question 3: In which plane does the point (4, -3, 0) lie?**Solution:**

Given: x, y, z coordinates of the point are 4, -3, 0.

As the distance of point along the z-axis is 0, the plane in which the point lies is the xy-plane.

Question 4: In which octant does each of the given points lie?

(i) (-4, -1, -6)

(ii) (2, 3, -4)

(iii) (-6, 5, -1)

(iv) (4, -3, -2)

(v) (-1, -6, 5)

(vi) (4, 6, 8)

Solution:

The position of a point in a octant is signified by the signs of the x, y, z coordinates.

Signs of x, y, z coordinates in all the octants are as follow:

Number	Sign of X	Sign of Y	Sign of Z
I	+	+	+
II	-	+	+
III	-	-	+
IV	+	-	+
V	+	+	-
VI	-	+	-
VII	-	-	-
VIII	+	-	-

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry

Compare signs with given table and allocate octants to the points

(i) $(-4, -1, -6)$ lies in octant VII

(ii) $(2, 3, -4)$ lies in octant V

(iii) $(-6, 5, -1)$ lies in octant VI

(iv) $(4, -3, -2)$ lies in octant VIII

(v) $(-1, -6, 5)$ lies in octant III

(vi) $(4, 6, 8)$ lies in octant I

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry**Exercise 26B**

Page No: 787

Question 1: Find the distance between the points :

- (i) A(5, 1, 2) and B(4, 6, -1)
- (ii) P(1, -1, 3) and Q(2, 3, -5)
- (iii) R(1, -3, 4) and S(4, -2, -3)
- (iv) C(9, -12, -8) and the origin

Solution:

The distance between two points (x_1, y_1, z_1) and (x_2, y_2, z_2) is given by

$$\text{Distance} = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

- (i) A(5, 1, 2) and B(4, 6, -1)

Here $(x_1, y_1, z_1) = (5, 1, 2)$ and $(x_2, y_2, z_2) = (4, 6, -1)$

Now,

$$\begin{aligned}\text{Distance} &= \sqrt{(4 - 5)^2 + (6 - 1)^2 + (-1 - 2)^2} \\ &= \sqrt{(-1)^2 + (5)^2 + (-3)^2} \\ &= \sqrt{1 + 25 + 9} \\ &= \sqrt{35}\end{aligned}$$

Distance between A and B is $\sqrt{35}$ units

- (ii) P(1, -1, 3) and Q(2, 3, -5)

Here $(x_1, y_1, z_1) = (1, -1, 3)$ and $(x_2, y_2, z_2) = (2, 3, -5)$

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry

$$\begin{aligned}\text{Distance} &= \sqrt{(2-1)^2 + (3-(-1))^2 + (-5-3)^2} \\ &= \sqrt{(1)^2 + (4)^2 + (-8)^2} \\ &= \sqrt{1+16+64} \\ &= \sqrt{81} \\ &= 9\end{aligned}$$

Distance between P and Q is 9 units

(iii) R(1, -3, 4) and S(4, -2, -3)

Here $(x_1, y_1, z_1) = (1, -3, 4)$ and $(x_2, y_2, z_2) = (4, -2, -3)$

$$\begin{aligned}\text{Distance} &= \sqrt{(4-1)^2 + (-2-(-3))^2 + (-3-4)^2} \\ &= \sqrt{(3)^2 + (1)^2 + (-7)^2} \\ &= \sqrt{9+1+49} \\ &= \sqrt{59}\end{aligned}$$

Distance between R and S is $\sqrt{59}$ units

(iv) C(9, -12, -8) and the origin

Here $(x_1, y_1, z_1) = (9, -12, -8)$ and $(x_2, y_2, z_2) = (0, 0, 0)$

$$\begin{aligned}\text{Distance} &= \sqrt{(0-9)^2 + (0-(-12))^2 + (0-(-8))^2} \\ &= \sqrt{(-9)^2 + (12)^2 + (8)^2} \\ &= \sqrt{81+144+64} \\ &= \sqrt{289} \\ &= 17\end{aligned}$$

Distance between C and Origin is 17 units.

Question 2: Show that the points A(1, -1, -5), B(3, 1, 3) and C(9, 1, -3) are the vertices of an equilateral triangle.

Solution:

Given points are the vertices of an equilateral triangle if measure of all the sides are equal.

Apply distance formula:

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry

The distance between two points (x_1, y_1, z_1) and (x_2, y_2, z_2) is given by

$$\text{Distance} = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

Length AB:

$$= \sqrt{(3 - 1)^2 + (1 - (-1))^2 + (3 - (-5))^2}$$

$$= \sqrt{(2)^2 + (2)^2 + (8)^2}$$

$$= \sqrt{4 + 4 + 64}$$

$$= \sqrt{72}$$

$$= 6\sqrt{2}$$

Length BC:

$$= \sqrt{(9 - 3)^2 + (1 - 1)^2 + (-3 - 3)^2}$$

$$= \sqrt{(6)^2 + (0)^2 + (-6)^2}$$

$$= \sqrt{36 + 0 + 36}$$

$$= \sqrt{72}$$

$$= 6\sqrt{2}$$

Length AC:

$$= \sqrt{(9 - 1)^2 + (1 - (-1))^2 + (-3 - (-5))^2}$$

$$= \sqrt{(8)^2 + (2)^2 + (2)^2}$$

$$= \sqrt{64 + 4 + 4}$$

$$= \sqrt{72}$$

$$= 6\sqrt{2}$$

$$\Rightarrow AB = BC = AC$$

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry

Therefore, Points A, B, C are vertices of an equilateral triangle.

Question 3: Show that the points A(4, 6, -5), B(0, 2, 3) and C(-4, -4, -1) form the vertices of an isosceles triangle.

Solution:

Given points are the vertices of an isosceles triangle if measure of any two sides are equal.
Apply distance formula:

The distance between two points (x_1, y_1, z_1) and (x_2, y_2, z_2) is given by

$$\text{Distance} = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

Length AB:

$$= \sqrt{(0 - 4)^2 + (2 - 6)^2 + (3 - (-5))^2}$$

$$= \sqrt{(-4)^2 + (-4)^2 + (6)^2}$$

$$= \sqrt{16 + 16 + 36}$$

$$= \sqrt{68}$$

$$= 2\sqrt{17}$$

Length BC:

$$= \sqrt{(-4 - 0)^2 + (-4 - 2)^2 + (-1 - 3)^2}$$

$$= \sqrt{(-4)^2 + (-6)^2 + (-4)^2}$$

$$= \sqrt{16 + 36 + 16}$$

$$= \sqrt{68}$$

$$= 2\sqrt{17}$$

Length AC:

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry

$$\begin{aligned} &= \sqrt{(-4 - 4)^2 + (-4 - 6)^2 + (-1 - (-5))^2} \\ &= \sqrt{(-8)^2 + (-10)^2 + (2)^2} \\ &= \sqrt{64 + 100 + 4} \\ &= \sqrt{168} \end{aligned}$$

From above result, we have $AB = BC$

Therefore, vertices A, B, C forms an isosceles triangle.

Question 4: Show that the points A(0, 1, 2), B(2, -1, 3) and C(1, -3, 1) are the vertices of an isosceles right-angled triangle.

Solution:

Given points are the vertices of an isosceles right-angled triangle if sum of squares of two sides is equal to square of third side.

Apply distance formula:

The distance between two points (x_1, y_1, z_1) and (x_2, y_2, z_2) is given by

$$\text{Distance} = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

Length AB:

$$\begin{aligned} &= \sqrt{(2 - 0)^2 + (-1 - 1)^2 + (3 - 2)^2} \\ &= \sqrt{(2)^2 + (-2)^2 + (1)^2} \\ &= \sqrt{9} \end{aligned}$$

Length BC:

$$\begin{aligned} &= \sqrt{(1 - 2)^2 + (-3 + 1)^2 + (1 - 3)^2} \\ &= \sqrt{(-1)^2 + (-2)^2 + (-2)^2} \\ &= \sqrt{9} \end{aligned}$$

Length AC:

$$\begin{aligned} &= \sqrt{(1 - 0)^2 + (-3 - 1)^2 + (1 - 2)^2} \\ &= \sqrt{1 + 16 + 1} \\ &= \sqrt{18} \end{aligned}$$

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry

$$\text{Now, } AB^2 + BC^2 = 9 + 9$$

$$= 18$$

$$= AC^2$$

Therefore, vertices A, B, C are the vertices of an isosceles right-angled triangle.

Question 5: Show that the points A(1, 1, 1), B(-2, 4, 1), C(1, -5, 5) and D(2, 2, 5) are the vertices of a square.

Solution:

Points A(1, 1, 1), B(-2, 4, 1), C(1, -5, 5) and D(2, 2, 5) are the vertices of a square if all sides are of equal measure.

Apply distance formula, and find the distance between all the points:

$$AB = \sqrt{(-2 - 1)^2 + (4 - 1)^2 + (1 - 1)^2}$$

$$= \sqrt{9 + 9 + 0}$$

$$= \sqrt{18}$$

$$BC = \sqrt{(-1 + 2)^2 + (5 - 4)^2 + (5 - 1)^2}$$

$$= \sqrt{1 + 1 + 16}$$

$$= \sqrt{18}$$

$$AC = \sqrt{(-1 - 1)^2 + (5 - 1)^2 + (5 - 1)^2}$$

$$= \sqrt{4 + 16 + 16}$$

$$= \sqrt{36}$$

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry

$$\begin{aligned}CD &= \sqrt{(2+1)^2 + (2-5)^2 + (5-5)^2} \\&= \sqrt{9+9+0} \\&= \sqrt{18}\end{aligned}$$

$$\begin{aligned}AD &= \sqrt{(2-1)^2 + (2-1)^2 + (5-1)^2} \\&= \sqrt{1+1+16} \\&= \sqrt{18}\end{aligned}$$

$$\begin{aligned}BD &= \sqrt{(2+2)^2 + (2-4)^2 + (5-1)^2} \\&= \sqrt{16+4+16} \\&= \sqrt{36}\end{aligned}$$

From above results, $AB = BC = CD = AD$, and $AC = BD$

Hence vertices A, B, C, D form a square.

Question 6: Show that the points A(1, 2, 3), B(-1, -2, -1), C(2, 3, 2) and D(4, 7, 6) are the vertices of a parallelogram. Show that ABCD is not a rectangle.

Solution:

Apply distance formula, and find the distance between all the points:

$$\begin{aligned}AB &= \sqrt{(-1-1)^2 + (-2-2)^2 + (-1-3)^2} \\&= \sqrt{4+16+16} \\&= \sqrt{36}\end{aligned}$$

$$\begin{aligned}BC &= \sqrt{(2+1)^2 + (3+2)^2 + (2+1)^2} \\&= \sqrt{9+25+9} \\&= \sqrt{43}\end{aligned}$$

$$\begin{aligned}CD &= \sqrt{(4-2)^2 + (7-3)^2 + (6-2)^2} \\&= \sqrt{4+16+16} \\&= \sqrt{36}\end{aligned}$$

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry

$$\begin{aligned}AD &= \sqrt{(4-1)^2 + (7-2)^2 + (6-3)^2} \\&= \sqrt{9 + 25 + 9} \\&= \sqrt{43}\end{aligned}$$

$$\begin{aligned}BD &= \sqrt{(4+1)^2 + (7+2)^2 + (6+1)^2} \\&= \sqrt{25 + 81 + 49} \\&= \sqrt{155}\end{aligned}$$

$$\begin{aligned}AC &= \sqrt{(2-1)^2 + (3-2)^2 + (2-3)^2} \\&= \sqrt{1 + 1 + 1} \\&= \sqrt{3}\end{aligned}$$

From above result, we have

$AB = CD$ [opposite sides]

$BC = AD$ [opposite sides]

Diagonal $AC \neq BD$, Diagonals are not equal

Hence, Points are vertices of parallelogram and ABCD is not a rectangle.

Question 7: Show that the points $P(2, 3, 5)$, $Q(-4, 7, -7)$, $R(-2, 1, -10)$ and $S(4, -3, 2)$ are the vertices of a rectangle.

Solution:

PQRS is a rectangle if opposite sides are equal and diagonals are also of equal length.

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry

$$\begin{aligned}PQ &= \sqrt{(-4-2)^2 + (7-3)^2 + (-7-5)^2} \\&= \sqrt{36 + 16 + 144} \\&= \sqrt{196}\end{aligned}$$

$$\begin{aligned}QR &= \sqrt{(-2+4)^2 + (1-7)^2 + (-10+7)^2} \\&= \sqrt{4 + 36 + 9} \\&= \sqrt{49}\end{aligned}$$

$$\begin{aligned}RS &= \sqrt{(4+2)^2 + (-3-1)^2 + (2+10)^2} \\&= \sqrt{36 + 16 + 144} \\&= \sqrt{196}\end{aligned}$$

$$\begin{aligned}PS &= \sqrt{(4-2)^2 + (-3-3)^2 + (2-5)^2} \\&= \sqrt{4 + 36 + 9} \\&= \sqrt{49}\end{aligned}$$

$$\begin{aligned}QS &= \sqrt{(4+4)^2 + (-3-7)^2 + (2+7)^2} \\&= \sqrt{64 + 100 + 81} \\&= \sqrt{245}\end{aligned}$$

$$\begin{aligned}PR &= \sqrt{(-2-2)^2 + (1-3)^2 + (-10-5)^2} \\&= \sqrt{16 + 4 + 225} \\&= \sqrt{245}\end{aligned}$$

Here, $PQ = RS$ and $QR = PS$ and $PR = QS$

Hence, points are vertices of a rectangle.

Question 8: Show that the points $P(1, 3, 4)$, $Q(-1, 6, 10)$, $R(-7, 4, 7)$ and $S(-5, 1, 1)$ are the vertices of a rhombus.

Solution:

Apply distance formula and find the length of each side.

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry

$$\begin{aligned}PQ &= \sqrt{(-1-1)^2 + (6-3)^2 + (10-4)^2} \\&= \sqrt{4+9+36} \\&= \sqrt{49}\end{aligned}$$

$$\begin{aligned}QR &= \sqrt{(-7+1)^2 + (4-6)^2 + (7-10)^2} \\&= \sqrt{36+34+9} \\&= \sqrt{49}\end{aligned}$$

$$\begin{aligned}RS &= \sqrt{(-5+7)^2 + (1-4)^2 + (1-7)^2} \\&= \sqrt{4+9+36} \\&= \sqrt{49}\end{aligned}$$

$$\begin{aligned}PS &= \sqrt{(-5-1)^2 + (1-3)^2 + (1-4)^2} \\&= \sqrt{36+4+9} \\&= \sqrt{49}\end{aligned}$$

$$\begin{aligned}QS &= \sqrt{(-5+1)^2 + (1-6)^2 + (1-10)^2} \\&= \sqrt{16+25+81} \\&= \sqrt{122}\end{aligned}$$

$$\begin{aligned}PR &= \sqrt{(-7-1)^2 + (4-3)^2 + (7-4)^2} \\&= \sqrt{64+1+9} \\&= \sqrt{74}\end{aligned}$$

$$\Rightarrow PQ = RS = QR = PS$$

And, diagonals: $PR \neq QS$

Hence, points are vertices of a rhombus.

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry

Exercise 26C

Page No: 793

Question 1: Find the coordinates of the point which divides the join of A(3, 2, 5) and B(-4, 2, -2) in the ratio 4 : 3.

Solution:

Section formula: The coordinates of point R that divides the line segment joining points A(x_1, y_1, z_1) and B(x_2, y_2, z_2) in the ratio m: n are

$$\left(\frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n}, \frac{mz_2 + nz_1}{m+n} \right)$$

Here, Point A(3, 2, 5) and B(-4, 2, -2),
m = 4 and n = 3

Using the above formula, we get,

$$= \left(\frac{4 \times -4 + 3 \times 3}{4+3}, \frac{4 \times 2 + 3 \times 2}{4+3}, \frac{4 \times -2 + 3 \times 5}{4+3} \right)$$

$$= \left(\frac{-7}{7}, \frac{14}{7}, \frac{7}{7} \right)$$

$$= (-1, 2, 1)$$

(-1, 2, 1) is the coordinates of the point which divides AB in the ratio 4 : 3.

Question 2: Let A(2, 1, -3) and B(5, -8, 3) be two given points. Find the coordinates of the point of trisection of the segment AB.

Solution:

Given: Point A(2, 1, -3) and B(5, -8, 3)
m = 2 and n = 1

Using section formula:

$$= \left(\frac{2 \times 5 + 1 \times 2}{2+1}, \frac{2 \times -8 + 1 \times 1}{2+1}, \frac{2 \times 3 + 1 \times -3}{2+1} \right)$$

$$= \left(\frac{12}{3}, \frac{-15}{3}, \frac{3}{3} \right)$$

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry

$$= (4, -5, 1)$$

Therefore, (4, -5, 1), is the point of trisection of the segment AB.

Question 3: Find the coordinates of the point that divides the join of A(-2, 4, 7) and B(3, -5, 8) externally in the ratio 2 : 1.

Solution:

Given: Point A(-2, 4, 7) and B(3, -5, 8)

$$m = 2 \text{ and } n = 1$$

Using section formula:

$$\left(\frac{2 \times 3 - 1 \times -2}{2 - 1}, \frac{2 \times -5 - 1 \times 4}{2 - 1}, \frac{2 \times 8 - 1 \times 7}{2 - 1} \right)$$

$$= (8, -14, 9),$$

Therefore, (8, -14, 9), is the point that divides AB externally in the ratio 2:1.

Question 4: Find the ratio in which the point R(5, 4, -6) divides the join of P(3, 2, -4) and Q(9, 8, -10).

Solution:

Given: Point P(3, 2, -4) and Q(9, 8, -10) divided by a point R(5, 4, -6)

Let the ratio be k:1 in which point R divides point P and point Q.

We know below are the coordinates of a point which divides a line segment in ratio, k:1.

Using section formula, we have

$$(5, 4, -6) = \left(\frac{k \times 9 + 1 \times 3}{k + 1}, \frac{k \times 8 + 1 \times 2}{k + 1}, \frac{k \times -10 + 1 \times -4}{k + 1} \right)$$

Find the value of k:

Choosing first co-ordinate,

$$5 = (9k+3)/(k+1)$$

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry

$$5k + 5 = 9k + 3$$

$$4k = 2$$

Therefore, the required ratio be 1:2.

Question 5: Find the ratio in which the point C(5, 9, -14) divides the join of A(2, -3, 4) and B(3, 1, -2).

Solution:

Given: Point A(2, -3, 4) and B(3, 1, -2) divided by a point C(5, 9, -14)

Let the ratio be k:1 in which point C divides point A and point B.

We know below are the coordinates of a point which divides a line segment in ratio, k:1.

Using section formula, We have

$$(5, 9, -14) = \left(\frac{k \times 3 + 1 \times 2}{k + 1}, \frac{k \times 1 + 1 \times -3}{k + 1}, \frac{k \times -2 + 1 \times 4}{k + 1} \right)$$

Find the value of k:

Choosing first co-ordinate,

$$5 = (3k+2)/(k+1)$$

$$5k + 5 = 3k + 2$$

$$2k = -3$$

$$\text{Or } k = -3/2$$

Since, the ratio is -3:2, division is external. Therefore, external division ratio is 3:2.

Question 6: Find the ratio in which the line segment having the end points A(-1, -3, 4) and B(4, 2, -1) is divided by the xz-plane. Also, find the coordinates of the point of division.

R S Aggarwal Solutions for Class 11 Maths Chapter 26 Three Dimensional Geometry

Solution: Let the plane XZ divides the points A(-1, -3, 4) and B(4, 2, -1) in ratio k:1.

Using section formula, we have

$$(x, 0, z) = \left(\frac{k \times 4 + 1 \times -1}{k + 1}, \frac{k \times 2 + 1 \times -3}{k + 1}, \frac{k \times -1 + 1 \times 4}{k + 1} \right)$$

Since we are given with xz-plane, so y-coordinate is zero here.

Choosing y-coordinate, we have

$$0 = (2k-3)/(k+1)$$

$$\text{or } k = 3/2$$

The ratio is 3:2 in XZ plane which divides AB.